



University
of Glasgow

Yaro, M., Munyard, K. A., Stear, M. J., and Groth, D. M. (2016) Molecular identification of livestock breeds: a tool for modern conservation biology. *Biological Reviews*, (doi:10.1111/brv.12265).

There may be differences between this version and the published version. You are advised to consult the publisher's version if you wish to cite from it.

<http://eprints.gla.ac.uk/124472/>

Deposited on: 14 September 2016

Enlighten – Research publications by members of the University of Glasgow
<http://eprints.gla.ac.uk>

Molecular Identification of Livestock Breeds: A Tool for Modern Conservation Biology

M Yaro¹, K.A. Munyard¹, M. J. Stear², D.M. Groth^{1*}

¹ School of Biomedical Sciences, CHIRI Biosciences Research Precinct, Faculty of Health Sciences, Curtin University, GPO Box U1987, Perth, WA, Australia 6845.

² Department of Animal Production and Public Health, Faculty of Veterinary Medicine, University of Glasgow, Bearsden Road, Glasgow G61 1QH.

* Corresponding author at: School of Biomedical Sciences, Curtin University, GPO Box U1987, Perth, Western Australia 6845, Australia. (Tel.: +618 92667475; fax: +618 92662342. E-mail address: D.Groth@curtin.edu.au).

ABSTRACT

Global livestock genetic diversity includes all of the species, breeds and strains of domestic animals, and their variations. Although a recent census had indicated that there were 40 species and over 8000 breeds of domestic animals; for the purpose of conservation biology the diversity between and within breeds rather than species is regarded to be of crucial importance. This domestic animal genetic diversity has developed through three main evolutionary events, from speciation (about 3 million years ago) through domestication (about 12,000 years ago) to specialised breeding (starting about 200 years ago). These events and their impacts on global animal genetic resources have been well documented in the literature. The key importance of global domestic animal resources in terms of economic, scientific and cultural heritage has also been quantified. In spite of the importance, there have been a growing number of reports on the alarming erosion of domestic animal genetic

resources. This erosion of animal genetic resources is happening in spite of several global conservation initiatives designed to mitigate it. In this review, we discuss these conservation interventions and highlight their strengths and weaknesses. However, pivotal to the success of these conservation initiatives is the reliability of the genetic assignment of individual members within a target breed. Finally, we discuss the prospect of using improved breed identification methodologies to develop a reliable breed specific molecular identification tool that is easily applicable to populations of livestock breeds in various ecosystems. These identification tools, when developed, will not only facilitate the regular monitoring of threatened or endangered breed populations, but also enhance the development of more efficient and sustainable livestock production systems.

Key words: Conservation, Diversity, Genetic Resources, Global Livestock, FAO, Molecular Techniques, Threats, Breeds.

CONTENTS

I. Introduction	3
II. Key Scientific, Cultural and Economic Importance of Global livestock Genetic Resources.....	6
III. Threats to Global livestock Genetic Resources.....	7
IV. Assessment of Livestock Genetic Diversity and Conservation Status.....	11
1. Variable Number of Tandem Repeats (VNTRs).....	14
2. Single Nucleotide Polymorphisms (SNPs) Marker.....	16
V. Conservation Methods for Domestic Animal Genetic Resources.....	17
VI. Modelling: The Way forward.....	21
VII. Conclusions.....	23

VIII. Acknowledgements.....	24
IX. References.....	24

I. INTRODUCTION

A significant amount of genetic variation present in wild animal lineages prior to domestication was conserved during the domestication process, and persisted within their respective domesticates (Dobney & Larson, 2006). Over the 12,000 years since farm animals were first domesticated, their genetic make-up has undergone adaptation due to both natural (speciation) and artificial (domestication/ breeding) selection pressures exerted by their specific environments and human activity respectively (Banik, Pankaj & Naskar, 2015; Hoffmann & Scherf, 2005; Jensen, 2006; Mignon-Grasteau, Boissy, Bouix *et al.*, 2005; Morris, 2006; Naskar, Gowane & Chopra, 2015; Price, 1999; Vigne, 2011; Zeder, Emshwiller, Smith *et al.*, 2006). These selection pressures have culminated in the development of a rich global domestic animal diversity with thousands of breeds (Ajmone-Marsan & Globaldiv Consortium, 2010; Groeneveld, Lenstra, Eding *et al.*, 2010). Each of these breeds is characterised by their unique morphology and productivity related to specific environmental and applied farming conditions (Lopes, Mendonça, Rojer *et al.*, 2015; Shand, 1997). A livestock breed can be generally defined as either a homogenous group with unique and identifiable phenotypic features that distinguish it from other sub groups within the same species, or a homogenous group for which geographical isolation from other groups of the same species has resulted in their acceptance as unique entities (FAO, 2000; Rege, 2003). Recently a more refined definition of a breed concept to encompass the history of the livestock was proposed by Feliuss, Theunissen and Lenstra (2014) and Tixier-Boichard (2014). This proposal will conform to current practical reality as not all breeds by definition actually represent unique genetic resources. Breeds can therefore be regarded as the unit

of management for livestock instead of unit of conservation so as to make it more instrumental for conservation purposes (Feliuss *et al.*, 2014; Groeneveld *et al.*, 2010).

A recent report on livestock breed diversity stated that there were 7202 local breeds (breeds found in only one country), 509 regional trans boundary breeds (breeds found in different countries within one region) and 551 international trans boundary breeds (breeds found in different countries in different continents) (FAO, 2013). These breed classifications comprised the seven main mammalian livestock species (sheep, goats, cattle, pigs, buffalo, horses, and asses/donkeys), four main avian livestock species (chicken, turkeys, ducks, and geese) and eight minor livestock species (alpacas, yaks, llamas, camels, elephants, musk oxen, and guinea pigs). However, since the concept of selective breeding emerged in the last 200 years, and subsequently, through more intensive selection in the last few decades, domestic animal diversity has been under sustained threat of significant erosion (Ajmone-Marsan *et al.*, 2010; Köhler-Rollefson, 1997). In 2012, an analysis of data from 182 countries at the Global Databank for farm animal genetic resources revealed that 8% of farm animal breeds could already be considered extinct, 22% were at varying degrees of extinction risk, and the risk status of 34% was unknown (FAO, 2013). The report indicated that only approximately 36% of global farm animal genetic resources were not at any risk of immediate extinction.

This growing threat to the world's animal genetic resources was recognised by the Food and Agriculture Organisation of the United Nations (UN) as an emerging global challenge, and this recognition has led to the ratification by 109 countries, in 2007, of the Interlaken Declaration on world animal genetic resources (Rischkowsky, Pilling, Food *et al.*, 2007). The Interlaken Declaration was the first ever such global action plan specifically aimed at conserving our current animal genetic resources. The declaration called for urgent and prompt measures to be undertaken to mitigate the risk of large scale loss of defined breeds in the face of challenges such as increasing human population, climate change and emerging diseases. It was also envisaged that such intervention,

when successful, would also make a significant contribution to millennium development goals 1 and 7:- eradication of extreme poverty and hunger, and ensuring environmental sustainability, respectively. The millennium development goals (or agenda) are a blue-print of eight goals referred to as the UN Millennium declaration, which was commissioned by the UN general assembly in September, 2000 (United Nations, 2000). The objective of the declaration is to galvanize unprecedented efforts from all member countries to reverse the poverty, hunger and disease affecting billions of people around the world within a 15 year time frame. Despite the historic breakthrough at the Interlaken Summit, not much progress has been made so far, especially in developing countries, due to several factors, the most prominent being a general lack of technical capacity and financial resources (FAO, 2007). The Domestic animal diversity information system (DAD-IS) is an information and communication tool that was set up to coordinate management strategies developed for domestic animal diversity at global, regional and national levels. This system has challenges, especially regarding the quality of the entries from developing countries (Tixier-Boichard, 2014). Most of the data submitted, especially from Africa, requires regular updating to make it relevant to the current situation. For example 48% and 53% of mammalian and avian breeds recorded in DAD-IS were found to lack sufficient demographic information necessary for the assessment of their respective risk status. Furthermore, 87% of entries regarding breed demographics were found to be based on a survey or census, thus presenting a significant limitation and might be unreliable. In recognition of these and other shortcomings in attempts at addressing the global animal genetic resource erosion issues, the European Union commissioned another three year global programme named 'The GLOBALDIV Project' (Ajmone-Marsan *et al.*, 2010). The GLOBALDIV project also known as "global view of livestock biodiversity and conservation" had representations from the FAO of UN, the International Livestock Research Institute (ILRI), the International Atomic Energy Agency (IAEA), and 34 individual international researchers from key institutions that are working in areas related to the characterisation of farm animal genetic resources. The main aim of this project is to integrate and disseminate the experience of past, large

scale, biodiversity projects and to review the main drivers of biodiversity loss, and to implement strategies for the conservation of farm animal genetic diversity. Notable among the recommendations of the GLOBADIV project is the need for amalgamation of the disciplines of genetics, socioeconomics and geographic information science for efficient valuation of domestic animal genetic resources. Currently, improved geo-referencing methods, for example global positioning systems (GPS), are being used as part of a range of measures to provide better production environment descriptors (Groeneveld *et al.*, 2010). However, because of the dynamic nature of domestic livestock diversity, it is now obvious that more innovative interventions are required to provide precise information on breed structure and status and effectively halt the rapid loss of global livestock genetic diversity. For any livestock breed considered to be at risk, it is recommended that the monitoring of population trends in terms of population size and structure must be carried out at least once per generation. The development of breed specific identification tools for each characterised livestock breed will not only facilitate this process of regular monitoring of population trends and demographics, but also promote conservation.

This review summarizes our knowledge of; (i) the key importance of domestic animal genetic resources, (ii) the threats to this resource diversity, (iii) the current status of domestic animal genetic resources, and (iv) conservation methods, with specific emphasis on a molecular genetics approach. We conclude with an assessment of the potential development and use of reliable breed identification tools for livestock breeds for enhancing modern conservation biology studies and preservation of livestock breed diversity.

II. KEY SCIENTIFIC, CULTURAL AND ECONOMIC IMPORTANCE OF GLOBAL LIVESTOCK GENETIC RESOURCES

Domestic livestock are known to directly provide food and livelihoods to more than 90% of the 1.97 billion people who live on less than one US Dollar a day (Anderson, 2003; ILRI, 2009). With a total global asset value of US\$ 1.4 trillion dollars, domestic livestock is reported to contribute 33% and

17% to global protein and kilocalorie consumption, respectively (Herrero, Thornton, Gerber *et al.*, 2009). In many developing countries, apart from the provision of food and income, livestock transactions also develop and foster meaningful and emotional social relationships between and among communities (McCorkle & James, 1996). The so called minor livestock species, although fewer in population number and distribution, are typically of critical importance in terms of cultural heritage and for the livelihood of their owners (McCorkle *et al.*, 1996; York & Mancus, 2013). For instance draught animal power plays essential role in the livelihoods of marginal communities in many developing countries in Asia, sub Saharan Africa, and Latin America (Barrett, 1992; Lawrence & Pearson, 2002; Teweldmehidin & Conroy, 2010). In addition to these traditional important uses, several species of animals are now used as models in toxicology studies to ascertain the hazard level to human of prospective drugs (Olson, Betton, Robinson *et al.*, 2000). For example the miniature pig was identified as the ideal non-human primate model for chromosomal abnormalities, skin cell therapy and neural stem cell studies (Vodička, Smetana, Dvořánková *et al.*, 2005). Also a strain of rabbit referred to as Watanabe heritable hyperlipidemic was found to be a good model for the study of human myocardial infarction (Shiomi, Ito, Yamada *et al.*, 2003). It has been recommended that comparative medicine, which entails disease studies across animals and human species, holds the key to efficient prevention and control strategies for many zoonotic diseases (Kahn, 2006). Livestock diversity should not only be considered on the basis of global food security, but also as having critical cultural, economic and scientific importance, both currently and into the future.

III. THREATS TO GLOBAL LIVESTOCK GENETIC RESOURCES

The global domestic animal or livestock genetic resources (AnGR) are defined as the sum total of animal species, breeds and strains that currently are, or may be of, future economic, scientific and cultural heritage importance to humans. For the purpose of conservation it is usually breed diversity rather than species diversity that is of greater importance (Philipsson, Zonabend, Bett *et al.*, 2011). According to the latest report by the commission on animal genetic resources the percentage of

local livestock breeds considered to be at the risk of extinction increased by two percentage points between 2010 and 2012 (FAO, 2013). This outlook on the prevailing extinction rate of livestock, although alarming, is likely to be an under-estimation of the actual situation, especially in relation to estimates for developing regions of the world such as sub-Saharan Africa (Rege & Gibson, 2003). The loss of livestock genetic diversity reduces the range of opportunities available to confront the challenges of unpredictable future events, such as climate change, social change, disease epidemics, selection errors, and many others (Anderson, 2003; Anderson & Centonze, 2007).

There is a wide spectrum of interrelated man-made and natural factors that pose varying levels of threats to global AnGR (Philipsson *et al.*, 2011; Rege *et al.*, 2003). The factors that are responsible for the erosion of genetic diversity are often a function of the size of the population under consideration (Barbato, Orozco-terWengel, Tapio *et al.*, 2015). Generally, the smaller a livestock population, the greater is its vulnerability to extinction (Biscarini, Nicolazzi, Stella *et al.*, 2015; Henson, 1992; Ramstad, Woody, Sage *et al.*, 2004). Human factors offer the greatest threat to global livestock diversity (Biscarini *et al.*, 2015; Frankham, 1995). The human factors include, but are not limited to; intensive selective breeding, over exploitation, political instability and wars (Goe & Stranzinger, 2002), indiscriminate crossbreeding (Alvarez, Traoré, Tamboura *et al.*, 2009; Wollny, 2003) and general neglect or lack of breeding programmes (Rege, Marshall, Notenbaert *et al.*, 2011; Wollny, 2003). Interestingly, these human factors vary across both developed and developing regions of the world. In the developed regions, the threat to livestock diversity is mostly associated with overexploitation such as specialised breeding in response to dynamic socioeconomic pressures (Groeneveld *et al.*, 2010). This trend is also expedited partially by often misguided or inappropriate application of advanced molecular biology technologies (Tisdell, 2003). Conversely in the developing countries, the main factors are a general neglect of livestock and or poorly structured breeding programmes driven in part by lack of technical knowledge and financial resources (Alvarez *et al.*, 2009; Biscarini *et al.*, 2015; Philipsson *et al.*, 2011). In the face of this clear dimorphism, what is of utmost importance are the measures necessary to minimise the “Swanson dominance effect” from

happening (Tisdell, 2003). The Swanson dominance effect refers to a phenomenon in which the choices made by the earliest developing societies influence the later pattern of development in another. There have been reports of livestock keepers in parts sub-Saharan Africa abandoning their locally adapted breeds in favour of specialised potentially highly productive, but non-adapted exotic breeds, thereby leading to a decline in diversity (Groeneveld *et al.*, 2010; Wollny, 2003). Nonetheless, regardless of the region of the world, general increases in human population tend to impact negatively on livestock diversity.

Natural events that have commonly been cited as major causes of erosion of livestock genetic resources include; Tsunamis, Earthquakes, Hurricanes, droughts, disease epidemics, famine and floods (Prentice & Anzar, 2011). In the past two or more decades, climate change has emerged as a higher-level driving force for reduction in AnGR (Nardone, Ronchi, Lacetera *et al.*, 2010; Thornton, van de Steeg, Notenbaert *et al.*, 2009). Many reports have described the expected impact of climate change on livestock production systems and diversity (Banik *et al.*, 2015; Herrero *et al.*, 2009; Hoffmann, 2010; Kantanen, Løvendahl, Strandberg *et al.*, 2015; McMichael, Powles, Butler *et al.*, 2007; Naskar *et al.*, 2015). This is mainly because of the direct and indirect implication of climate change on both the frequencies and intensities of most of the causative factors for genetic erosion mentioned previously (Naskar *et al.*, 2015). The irony, however, is that livestock contributes significantly to climate change, as they contribute about a fifth of global greenhouse gas emissions (Garnett, 2009; Gavrilova, Jonas, Erb *et al.*, 2010; McMichael *et al.*, 2007; Shields & Orme-Evans, 2015).

Natural and human-made evolutionary forces either directly or indirectly can cause a reduction in the effective population size (N_e) of a livestock breeding population. Therefore, the genetic variability of subsequent populations is drastically reduced because it is derived from the genetic constitution of the few survivors remaining from the original population (Allendorf, 1986). In population genetic studies these reductions in population size are referred to bottlenecks. A

population that passes through a bottleneck loses alleles and usually shows reduced average heterozygosity (Allendorf, 1986; Nei, Maruyama & Chakraborty, 1975), but could also lead to temporarily, an increased in heterozygosity if more rare alleles are lost in the process (Hundertmark & Van Daele, 2010; Luikart & Cornuet, 1998). Regardless of the cause of a bottleneck, it may take several generations to restore the original level of heterozygosity through the occurrence of new mutations (Chakraborty & Nei, 1977). The impact of a bottleneck is logically more profound on small breeding populations because the originally available pool of genetic diversity is smaller, and is hence more severely depleted in fewer generations. In population genetic studies a bottleneck effect is referred to as a founder effect if it is associated with the founding of a new population (Dlugosch & Parker, 2008; Ramstad *et al.*, 2004; Templeton, 1980). Random events such as founder and bottleneck effects that imperfectly eliminate genes and reduce variability within a population are described as genetic drift (Newman & Pilson, 1997; Ramstad *et al.*, 2004). Reduction in heterozygosity in a livestock population can be associated with decline in fitness of individual members, as is often the case in wild populations (Worley, Collet, Spurgin *et al.*, 2010). This is because within small populations, the rate of inbreeding is much higher and consequently there is higher likelihood of the expression of deleterious recessives in a homozygous state. The expression of deleterious alleles has adverse effects on the livestock population often presenting as reduced production, reproduction and survival (Dlugosch *et al.*, 2008; Lacy, 1997). Frankham (1995) and Lacy (1997) have described the positive correlation between inbreeding and risk of extinction. The effective population size model takes into account important population variables such as age and structure, inbreeding rates, genetic drift, genetic diversity and sex ratio. For example, a population of four males and four females constitutes the same effective population size as that of 100 females and only two males (Henson, 1992).

In a breed regeneration programme, the effective population size can be enhanced by equalizing the male to female ratio, and standardizing litter size and longevity within the breeding population, so as to ensure that each animal contributes equally to the next generation. Therefore the effective

population size is the preferred indicator of livestock conservation risk status (Dlugosch *et al.*, 2008; Nei *et al.*, 1975). However, it is apparent that the estimation of the effective population size and, subsequent determination of its conservation status for a given breed is limited by the lack of availability of a reliable breed identification tool for any specific breed.

IV. ASSESSMENT OF LIVESTOCK GENETIC DIVERSITY AND CONSERVATION STATUS

In order to sustainably manage livestock genetic resources a comprehensive knowledge of diversity within and between breed populations is required (Groeneveld *et al.*, 2010). A major step towards standardising the assessment criteria for livestock breed conservation status was the establishment of a universal classification framework by the FAO for categorizing risk status. The current classification of livestock conservation risk status contains seven categories, namely; extinct, critical, critical-maintained, endangered, endangered-maintained, not at risk, and unknown (FAO, 2013). Regular assessment of genetic conservation status of livestock is of fundamental importance to prevent genetic erosion and to preserve diversity. Key to achieving an effective assessment of livestock conservation status is a reliable mode of identification of members of a target breed. There are two broad methods for identifying individual members of a livestock breed, and their merits and demerits have been discussed thoroughly (Agaviezor, Peters, Adefenwa *et al.*, 2012; Ashley & Dow, 1994; Birteeb, Peters, Yakubu *et al.*, 2012). These methods comprise phenotypic and molecular identification techniques. Traditionally phenotypic identification has been used to identify the breed of an individual in livestock genetic diversity studies. The phenotypic parameters usually used comprise; physical features (e.g. Shape of horn, ears, body measurements, colour etc.), production traits (e.g. growth parameters), reproductive traits (e.g. fecundity) and survival traits (e.g. Disease resistance, drought resistance) (Brinks, Clark, Kieffer *et al.*, 1964; Gwakisa, Kemp & Teale, 1994; Reverter, Johnston, Ferguson *et al.*, 2003). These methods are extensively used because they are inexpensive and often do not require the use of any sophisticated equipment. However, the major

disadvantage is that the genetic diversity is observed only at the phenotypic level and this does not always correspond to actual diversity at the DNA level (Feliu *et al.*, 2014).

It is possible to find different phenotypes with similar genotypes, typically due to genotype-environment interactions, for example as observed in Brazilian sheep Breeds (Paiva, Faria, Silvério *et al.*, 2005) and Egyptian sheep breeds (Ali, 2003). Similar phenotypes with different genotypes also occur, as observed in West African Djallonke sheep (Alvarez, Capote, Traoré *et al.*, 2012; Alvarez *et al.*, 2009; Wafula, Jianlin, Sangare *et al.*, 2005). As a result, the use of molecular tools in many assessment studies of genetic diversity in different regions of the world revealed varying degrees of unexpected introgression and admixture in livestock populations. Some of the reported studies include; the Djallonke sheep breed of sub-Saharan Africa (Alvarez *et al.*, 2009; Wafula *et al.*, 2005), Herdwick sheep of the United Kingdom (Bowles, Carson & Isaac, 2014) and alpaca and llama of Latin America (Kadwell, Fernandez, Stanley *et al.*, 2001). This obvious shortcoming has rendered the use of the phenotypic method alone unreliable for determination of livestock breeds for the purpose of genetic diversity studies.

In livestock genetic diversity studies, the molecular method for determining breed identity entails two main approaches based upon either protein markers or DNA markers (Ferguson, Taggart, Prodöhl *et al.*, 1995; McMahon, Teeling & Höglund, 2014). Protein markers, also referred to as allozymes, are based on the characteristic polymorphism of the blood group systems, leucocyte antigens and enzymes (Dodgson, Cheng & Okimoto, 1997). This molecular method employs these protein markers to estimate genetic variability in livestock populations as well as phylogenetic relationships between breeds (Pepin & Nguyen, 1994; Witko-Sarsat, Friedlander, Capeillere-Blandin *et al.*, 1996). Although better than the phenotypic method, the use of protein markers is too expensive for large number of loci, and lacks the power to resolve differences between closely related breeds, because of limits of detection of genetic variation (Engel, Linn, Taylor *et al.*, 1996; Ferguson *et al.*, 1995; Toro, Fernández & Caballero, 2009). The use of DNA markers is the most

reliable molecular method for assessment of genetic diversity (Liu & Cordes, 2004). Nuclear and mitochondrial DNA marker analyses have revealed detailed information on many domestication events, such as their timing and location (Bruford, Bradley & Luikart, 2003; Zhao, Zheng, Dong *et al.*, 2013). DNA marker analyses provide an added opportunity for investigating the genetic composition of both extinct and endangered breeds without destructive sampling. There are seven principal DNA marker techniques commonly used for livestock diversity studies (Sunnucks, 2000). The seven DNA marker principal techniques have been discussed thoroughly and their advantages and disadvantages are well documented by many authors. These techniques are; Restriction Fragment length Polymorphism (RFLP) (Beckmann & Soller, 1983; Beckmann & Soller, 1986; Thurston, Siggins, Mileham *et al.*, 2002), Mitochondrial DNA barcoding (mtDNA) (Avisé, Arnold, Ball *et al.*, 1987; Avisé & Ellis, 1986; Harrison, 1989; Kocher, Thomas, Meyer *et al.*, 1989; Zhang & Hewitt, 1996), Random Amplified Polymorphic DNA (RAPD) (Ali, Huang, Qin *et al.*, 2004; Dodgson *et al.*, 1997; Koh, Lim, Chua *et al.*, 1998; Levin, Crittenden & Dodgson, 1993), Amplified fragment Length Polymorphism technique (AFLP) (Bleas, De Grandis, Lee *et al.*, 1998; Parsons & Shaw, 2001; Savelkoul, Aarts, De Haas *et al.*, 1999), Y-chromosome technique (Bruford *et al.*, 2003; Zeder *et al.*, 2006), and the two most popular DNA techniques are Variable Number of Tandem Repeats (Minisatellite and Microsatellite markers) (Chistiakov, Hellemans & Volckaert, 2006; Fan & Chu, 2007; Lopes *et al.*, 2015; Zane, Bargelloni & Patarnello, 2002) and Single Nucleotide Polymorphism (SNP) based techniques (Andersson & Georges, 2004; Liu *et al.*, 2004; McMahon *et al.*, 2014; Morin, Luikart & Wayne, 2004; Vignal, Milan, SanCristobal *et al.*, 2002).

The advancement of DNA technologies during the past three decades, and particularly since 2007 when high throughput next generation sequencing became readily available, is revolutionising livestock population genetics studies (Helyar, Hemmer-Hansen, Bekkevold *et al.*, 2011; Schlötterer, Kofler, Versace *et al.*, 2014). This revolution is expedited by the concomitant advancement in bioinformatics tools and pipelines (Kofler, Nolte & Schlötterer, 2015). DNA markers have been used not only for diversity studies but also for molecular characterisation of numerous livestock breeds

worldwide (Agaviezor *et al.*, 2012; Al-Atiyat, Salameh & Tabbaa, 2014; Alvarez *et al.*, 2012; Bowles *et al.*, 2014). The dramatic reduction in the cost of use of DNA markers has facilitated their greater use by researchers. For the purpose of this review the Variable Number Tandem Repeats (VNTR) and the Single Nucleotide Polymorphism (SNP) techniques will be discussed in greater detail because of their wider application compared with the other molecular markers types. AFLP and RAPD markers are both bi-allelic, but dominant in nature, and hence are less informative and also have low reproducibility compared to the other markers (Vignal *et al.*, 2002). These characteristics have rendered them less popular for most animal based molecular genetic studies. RFLP markers are bi-allelic and co-dominant, and were famously used in the first large scale mapping of the human genome. However, they have now been superseded by multi-allelic and more informative microsatellite markers in both animal and human genome studies which, in turn have been largely superseded by single nucleotide polymorphism (SNP) arrays. MtDNA along with microsatellite markers were once a popular molecular genetic techniques of choice for evolutionary and ecological studies, however the molecular information provided by mtDNA markers is limited to only maternally inherited loci (Morin *et al.*, 2004). The use of mtDNA techniques, in combination with archaeological data, have provided precise information on most of the important centres of domestication for the main livestock species around the world (Bruford *et al.*, 2003; GLOBALDIV CONSORTIUM, 2010; Guo, Savolainen, Su *et al.*, 2006; Zeder *et al.*, 2006). Similarly limited, the use of Y-chromosomal haplotype markers elucidates specific molecular information only on paternal inherited traits (Luikart, Fernández, Mashkour *et al.*, 2006).

IV 1. VARIABLE NUMBER OF TANDEM REPEATS (VNTRS)

The application of VNTRs for assessment of genetic variation, sub structuring and hybridization in natural populations has been well reviewed in great detail by many investigators (Bruford & Wayne, 1993; Chistiakov *et al.*, 2006; Fan *et al.*, 2007; Sunnucks, 2000). The VNTR technique is based on the abundance of tandem repeats of simple sequences of nucleotides throughout the eukaryotic

genome (Takezaki & Nei, 2008). These VNTRs have been categorised into minisatellites and microsatellites according to the number of nucleotides per motif of repeats. VNTRs of between 1 to 6 nucleotide base pair units are referred to as microsatellites (Ashley *et al.*, 1994; Chistiakov *et al.*, 2006; Fan *et al.*, 2007), whereas a range of between 10 to 60 nucleotide base pair units is regarded as a minisatellite (Ashley *et al.*, 1994; Wasko & Galetti, 2003). Whereas minisatellites are concentrated towards the telomere of chromosomes, microsatellites are randomly distributed in chromosomes. Microsatellite markers are highly polymorphic, codominant markers of relatively small size, and hence are more amenable to PCR typing than are minisatellites (Zane *et al.*, 2002). Also, in comparison to the RFLP and RAPD techniques, the genetic basis of variability is readily apparent for microsatellites. Most microsatellites are located in non-coding regions of the genomes, and very few are located within the coding regions (Chistiakov *et al.*, 2006). Generally, microsatellite primers developed for one species of livestock are broadly applicable to other closely related species. For example, microsatellite markers developed for studies in bovine species are applicable for caprine and ovine species (Engel *et al.*, 1996). The reason for this wide applicability of microsatellite markers stems from the fact that they are conserved across wide taxa of related species. This versatility has led to the popularity of microsatellite maps for economically important livestock species (Sunnucks, 2000).

Microsatellites have been used in linkage mapping in diverse organisms, for example in the bovine genome (Barendse, Vaiman, Kemp *et al.*, 1997), porcine genome (Rohrer, Alexander, Keele *et al.*, 1994), human genome (Dib, Fauré, Fizames *et al.*, 1996), and ovine genome (Maddox, Davies, Crawford *et al.*, 2001). Microsatellite have also been employed for the identification of Quantitative Trait Loci (QTL) in major livestock species, for example, carcass composition and growth rate in cattle (Casas, Shackelford, Keele *et al.*, 2000), back fat thickness and intramuscular fat in pigs (Rohrer & Keele, 1998) and intestinal parasitic infection in sheep (Davies, Stear, Benothman *et al.*, 2006). Other population genetics studies accomplished with microsatellite markers include; determination of evolutionary relationships (Buchanan, Adams, Littlejohn *et al.*, 1994), estimation of pedigree errors

(Visscher, Woolliams, Smith *et al.*, 2002), [determination of genetic diversity among livestock populations](#) (Alvarez *et al.*, 2012; Alvarez *et al.*, 2009; Marletta, Tupac-Yupanqui, Bordonaro *et al.*, 2006; Medugorac, Veit-Kensch, Ramljak *et al.*, 2011; Wafula *et al.*, 2005), and last but not least, genetic distance between livestock breeds (Alvarez *et al.*, 2012; Buchanan *et al.*, 1994; Vanhala, Tuiskula-Haavisto, Elo *et al.*, 1998). The genetic distance of individuals within a livestock population indicate the suitability of an individual for conservation purposes. Individuals within the same breed with the widest differences in genetic distances are deemed most suitable candidates for conservation programmes. The estimates of genetic distances are also relevant for the determination of divergence time and construction of phylogenies (Takezaki & Nei, 1996).

IV 2. SINGLE NUCLEOTIDE POLYMORPHISM (SNP) MARKERS

The growing importance of SNP marker applications in molecular genetics has been reviewed in detail by Barbato *et al.* (2015), Broxham (2015), Goddard and Hayes (2009), Vignal *et al.* (2002), Hamblin, Warburton and Buckler (2007) and Morin *et al.* (2004). Prior to the use of SNP markers, microsatellites were the most popular and efficient technique for genetic diversity investigation, not only in livestock but also in humans as well. SNPs represent a location within a DNA sequence for which more than one nucleotide type is present within a given population (Morin *et al.*, 2004). In a strict molecular sense, SNPs are base substitutions within nucleotide sequences, and the very high density of their occurrence in the genomes of eukaryotes, including livestock, has been of great significance in population genomics studies (Goddard *et al.*, 2009; Vignal *et al.*, 2002). Although SNPs are bi-allelic (sometimes tri-allelic or quadri-allelic), co-dominant molecular markers, their high density permits, more than any other technique, very detailed information to be elucidated on genome dynamics within a study population (Hamblin *et al.*, 2007; Morin *et al.*, 2004). Furthermore, they provide deeper insight than microsatellites with respect to linkage disequilibrium and haplotype diversity, pedigree information as well as past demographic events, such as bottlenecks within a target population (Clarke, Henry, Dodds *et al.*, 2014; Helyar *et al.*, 2011; Morin *et al.*, 2004;

Vignal *et al.*, 2002). These features of SNP markers, coupled with a relatively low error rate, are opening opportunities for wider applications of SNP markers in understanding of livestock genetic architecture, such as precise identification of genomic regions that control traits of economic and survival importance (Kohn, Murphy, Ostrander *et al.*, 2006; Matukumalli, Lawley, Schnabel *et al.*, 2009) and ultimately genomic selection (Clarke *et al.*, 2014; Goddard *et al.*, 2009). These advances in genetic marker application for use in population genetic studies will not only enhance the development of improved livestock production systems, but most importantly will facilitate the development of efficient conservation strategies. High density SNP chips have been used to accurately and correctly cluster sheep breeds from different continents (Kijas, Townley, Dalrymple *et al.*, 2009), and more recently for the effective management of highly endangered breeds of small cattle in the Balkans (Broxham, 2015). A study involving typing 47,594 snps from pigs derived from domestic and wild populations on the Iberian Peninsula showed that principal component analysis could assign animals to a specific population (Herrero-Medrano, Megens, Groenen *et al.*, 2013). Furthermore, effective population size, past effective population size and runs of homozygosity could be determined from the data. This shows that snp chips can be used to determine relationships between populations and levels of inbreeding.

V. CONSERVATION METHODS FOR DOMESTIC ANIMAL GENETIC RESOURCES

Conservation of AnGR basically comprises all the management practices carried out to preserve the pool of genetic diversity of livestock for the purpose of meeting the current and future needs of humans (Rege *et al.*, 2003). The relevance of conservation of AnGR has been thoroughly discussed by many authors, from several different perspectives, for example; economic evaluation as a basis for AnGR conservation decisions (Drucker, Gomez & Anderson, 2001), the role of cryopreservation, reproductive technologies and genetic resource banks for AnGR conservation strategies (Hiemstra, van der Lende & Woelders, 2006b; Holt & Pickard, 1999; Mara, Casu, Carta *et al.*, 2013), information on population kinships as a basis for AnGR conservation decisions (Eding & Meuwissen, 2001) and

the challenge of conserving indigenous AnGR diversity (Mendelsohn, 2003). Each breed of livestock consists of unique sets of genes resulting from evolutionary events and diverse selection pressures imposed by the environment combined with the activities of humans over time. It is therefore difficult, if not impossible, to replace lost breeds of livestock, because those unique evolutionary processes cannot be re-created. There has been a general consensus on three critical approaches regarding the conservation of domestic livestock breeds, and these are; sustainable utilization of available livestock breeds, appropriate diversity based improvement strategies for livestock breeds, and development of appropriate assessment and preservation strategies (FAO, 2000; Hammond, 1999; Koehler-Rollefson & Meyer, 2014; Notter, Mariante & Sheng, 1994; Thornton, Boone, Galvin *et al.*, 2007). In addition to these approaches, the FAO has also recommended the regular monitoring of livestock breed conservation status. Currently the two main methods of AnGR conservation applied are the *in situ* and the *ex situ* methods. The applicability of both conservation methods, and their respective merits and demerits has been reported extensively (Boettcher, Tixier-Boichard, Toro *et al.*, 2010; Hammond, 1994; Henson, 1992; Mara *et al.*, 2013; Rege *et al.*, 2003). *In situ* conservation can best be described as the sustainable breeding of an endangered livestock breed in their normal adaptive production environment, or as close to it as practically possible, to conserve genetic diversity over a long time (Andrabi & Maxwell, 2007; Henson, 1992). Notable features of *in situ* conservation therefore include selection and mating programmes that enhance genetic variation within the target group, as well as management of the ecosystem to sustain their production. The basic requirements for *in situ* conservation programmes are generally readily available and affordable globally. There is a distinct difference between developed and developing countries regarding the minimum number of individuals required to commence an *in situ* programme. This is typically due to the general differences in the efficiency of management of their respective livestock production systems. There are different minimum numbers of animals reported to be required for *in situ* conservation. For example, whereas the minimum number for major livestock breeds (i.e. cattle, sheep, goats, pigs) required for *in situ* conservation is 100-1000 breeding females in developed

countries, no fewer than 5000 breeding females is recommended for developing countries (Signorello & Pappalardo, 2003). Simon (1999) reported 500 breeding females for pigs and goats, 750 for cattle, and 1500 for sheep for European breeds. It has been recommended that, ideally, for unrelated animals a minimum of 25 males and 50 females is sufficient to commence an *in situ* conservation programme, because the possible loss of genetic variability is estimated to be less than 1% per generation (Henson, 1992; Mara *et al.*, 2013).

There are a number of flagships *in situ* conservation programmes in place to conserve and improve some disease resistant breeds of livestock in some African countries, for example, Ndama cattle in the republic of Guinea, Djallonke sheep in Ghana, and Tswana sheep in Botswana (Henson, 1992; Kosgey & Okeyo, 2007; Yapi-Gnaoré, Dagnogo & Oya, 2003). The unique advantage of the *in situ* conservation method is that the target livestock continues to be utilized in the process. However, the danger is that the target livestock remains susceptible to uncertain demographic threats such as natural disasters and disease epidemics. The *ex situ* livestock conservation method is the preservation of endangered livestock outside their normal production systems (Henson, 1992; Hiemstra, Drucker, Tvedt *et al.*, 2006a). This method is normally applied to target groups that are faced with imminent extinction, and hence requires the use of high level expertise and technology. The three main *ex situ* methods are cryopreservation, farm park conservation, and breed pools or composite preservation.

Cryopreservation, also referred to as *in vitro ex situ* is undoubtedly the most popular of the *ex situ* approaches to conservation of AnGR (Hiemstra *et al.*, 2006a). This approach involves the cryopreservation of eggs, semen and or embryos of endangered or threatened animals in genome banks for use in managing diversity or regenerating the population decades or even centuries later (Chen, Zhang & Yu, 2008; Hanks, 2001; Russo, Martelli, Mauro *et al.*, 2007; Xiao-Yong, Shu-Jun, Run-Zi *et al.*, 2008). Cryogenic storage of carefully evaluated genetic material from a target breed population is also seen as an insurance policy against future loss. The merits and demerits of using

these approaches have been previously discussed (Boettcher *et al.*, 2010; Munro & Adams, 1991; Philipsson, Rege & Okeyo, 2006; Pintado & Hourcade, 2011; Ruane & Sonnino, 2011). The application of cryopreservation formerly depended only on assisted reproductive techniques such as artificial insemination and embryo transfer technologies. However, recent advances in reproductive biotechnologies including semen sexing, embryo micromanipulation and *in vitro* fertilization have the potential to revolutionise the livestock cryopreservation approach (O'Brien, Steinman & Robeck, 2009; Prentice *et al.*, 2011). The cryopreserved genetic materials are shielded from the influence of the unfavourable environmental conditions in existence in the normal production ecosystems. Regeneration of a breed through only preserved semen requires a number of back crosses to be done. However, the entire genetic composition of the original breed is not recoverable with only cryopreserved semen (Andrabi *et al.*, 2007). In practice, the *in situ* and *ex situ* conservation methods are not mutually exclusive because the cryopreservation approach can be used to complement the *in situ* method to achieve better regeneration of endangered populations. A range of combinations of *in situ* and *ex situ* conservation methods are being applied in a now popular integrated conservation approach (de Souza, Batista, Melo *et al.*, 2011; Hiemstra *et al.*, 2006a). It has been recommended that a stock of cryopreserved semen from 25 unrelated sires is sufficient to provide a reasonable diversity to an endangered population (Bruns & Glodek, 1999; Mara *et al.*, 2013).

The farm park *ex situ* conservation approach is similar to the *in situ* conservation approach, except that the targeted breeds are preserved outside their normal production environment in a specialised institutional setting, also referred to as an Ark-farm (Simon, 1999). Farm park animals are usually also subjected to more stringent management regimes to conserve natural levels of genetic variability within each species (Chesser, Smith & Brisbin, 1980). A notable feature of the Farm Park approach is its popularity in attracting tourists, and hence creating awareness of the need to conserve endangered animals. The Cotswold farm park in the UK is an example where rare breeds of sheep, goats, cattle, pigs and horses are being conserved, and it attracts more than 100,000 visitors yearly (Henson, 1992). The breed pool preservation programme is unique in the sense that it

involves the breeding together of a pool of two to four rare breeds with similar characteristics, and subsequently managing their offspring to conserve genetic variation (Henson, 1992). It is, however, recommended that the breed characteristics of each of the rare breeds is well ascertained prior to commencing a breed pool programme (Santos, Affonso, Diniz *et al.*, 2013). This method is particularly suitable for genes that control obvious morphological traits and extreme quantitative traits such as coat colour and prolificacy, respectively. Although this approach conserves useful genes from the pool, individual breeds are lost in the process. An example of the breed pool approach is the conservation programme of four rare desert goat breeds, in the north eastern part of Brazil (Henson, 1992).

Given that no single conservation method is capable of solving the myriad of challenges of domestic animal genetic resource erosion, an integrated conservation approach has been advocated to provide more efficiency (Rege *et al.*, 2003).

VI. MODELLING: THE WAY FORWARD

The successful domestication of animals represents a pivotal historic event in the cultural and demographic development of humans. The importance of global domestic livestock diversity to human wellbeing is now well-appreciated. This is evident by the globally coordinated efforts directed at halting the decline in AnGR as well as the sustainable utilisation of available livestock resources as discussed in this review. These global initiatives have yielded several interventions which are being implemented at the international, regional and local level (Ajmone-Marsan *et al.*, 2010; FAO, 2013). The main global body is the DAD-IS that coordinates regional bodies, for example EFABIS (European farm Animal Biodiversity information System) and DAGRIS (Domestic Animal Genetic Resource information System) for European and African regions respectively. The regional bodies in turn coordinate the local or national bodies, which essentially are the individual member states of the FAO of the United Nations. These efforts are being supplemented by the activities of other important organisations, prominent members including; the GLOBALDIV, the International Society

for Animal Genetics (ISAG), the SAVE foundation and several livestock breed societies worldwide (Broxham, 2015). Some notable progress has been made towards reducing the rate of erosion of global AnGR. For example, the status information for global mammalian and avian livestock breeds in the DAD-IS has increased from 43% and 39% respectively in 2009, to 57% and 48% in 2012 respectively (FAO, 2013). The effective monitoring of breed conservation status of livestock requires at least one census per generation of that target breed (FAO, 2007; Groeneveld *et al.*, 2010). Hence, a specific breed identification tool for each livestock breed will expedite such an exercise. However, pivotal to the success of these conservation efforts is the reliability of genetic identification of individual members within a target breed. The advancement in molecular technology in the last two decades has significantly increased our understanding the population genetics of domestic animals. It is apparent that the molecular characterisation of all domestic livestock breeds particularly in developing countries is a pre-requisite for their sustainable utilisation and conservation. This is because characterisation at the molecular level provides precise information for the determination of the actual population characteristics such as genetic variation and the effective population size (Luikart, England, Tallmon *et al.*, 2003). Currently, many domestic livestock breeds, particularly those in the developing countries, have yet to be characterised due to myriad issues including the lack of financial and technological capacity, as already described previously in this review. A recent report indicated that the risk status of 36% of all populations of local livestock breeds still remains unknown (FAO, 2013).

The main molecular technique used for most livestock genetic characterisation was microsatellite markers, for example they were used in; Spanish native cattle breeds (Martín-Burriel, García-Muro & Zaragoza, 1999), Aberdeen Angus cattle breeds (Vasconcellos, Tambasco-Talhari, Pereira *et al.*, 2003), Austrian sheep breeds (Baumung, Cubric-Curik, Schwend *et al.*, 2006) and indigenous goats in sub-Saharan Africa (Chenyambuga, Hanotte, Hirbo *et al.*, 2004). Although highly informative, the current panels of microsatellites used for analyses are not capable of elucidating all the information required regarding breed variation in livestock (Toro *et al.*, 2009). Recently, it is

becoming more evident that SNP analysis is more suited for the high throughput genotyping that is required to elucidate greater molecular insights such as historical signatures of selection (Qanbari, Pausch, Jansen *et al.*, 2014), phenotypic variations within livestock breeds (Groenen, Megens, Zare *et al.*, 2011) as well as linkage disequilibrium over short physical distances (Kijas, Porto-Neto, Dominik *et al.*, 2014) than any of the other molecular markers currently available. The availability and accessibility of comprehensive databases of genomic data for various uses has also facilitated population genetic studies globally for example; the National Centre for Biotechnology Information (NCBI) (Sayers, Barrett, Benson *et al.*, 2011), Livestock Animal Quantitative Trait Loci (QTL) database (Hu, Park, Wu *et al.*, 2013), and University of California Santa Cruz (UCSC) genome browser (Dreszer, Karolchik, Zweig *et al.*, 2012). The challenge now is to use these enhanced insights and understanding of molecular methods to develop breed specific identification tools that are easily applicable to populations of livestock in different ecosystems. Such a breed-specific tool can be developed through identification and characterisation of unique phylogenomic SNPs in next generation sequenced pooled-genomic DNA from a minimum of 25 unrelated pure bred individuals. Information from any discovered phylogenomic SNPs can then be used to develop multiplexed SNP assays for the easy and precise identification of pure bred members from mixed populations of breeds in various ecosystems. These tools will not only facilitate the timely diagnosis of the conservation status of livestock breeds, but will also permit the regular monitoring of endangered breed populations, particularly in developing countries where the lack of technical and financial capacity is reported a major impediment.

VII. CONCLUSIONS

- (1) The importance of maintaining global domestic animal genetic diversity is important to human wellbeing.

- (2) Breed specific molecular identification tools are urgently needed so that reliable and expeditious identification of individual members of any given breed can be achieved, and this is a pre-requisite for sustainable utilization and conservation of any breed.
- (3) A growing number of studies have established that whole genome sequencing of pools of individuals within a group or breed provides great deal of information on genetic variation across the whole genome even when performed at relatively low coverage, but also at a considerable lower cost (Clarke *et al.*, 2014; Gautier, Foucaud, Gharbi *et al.*, 2013; Kim, Li, Guo *et al.*, 2010; Kofler *et al.*, 2015). This principle will be a cost effective technique for the identification of breed specific phylogenomic SNPs within a target breed for the purpose of developing a breed specific molecular identification tools.

VIII. Acknowledgements

We are grateful to CHIRI Biosciences and Curtin University of Technology for the use of its facilities. M.Y. was funded by grant from Curtin University International Strategic Research Scholarship (CIPRS).

IX. REFERENCES

- AGAVIEZOR B. O., PETERS, S. O., ADEFENWA, M. A., YAKUBU, A., AJAYI, O. O. & AMUSAN, S. A. (2012). Morphological and microsatellite DNA diversity of Nigerian indigenous sheep. *Journal of animal science and biotechnology* **3**, 38.
- AJMONE-MARSAN P. & GLOBALDIV CONSORTIUM, T. (2010). A global view of livestock biodiversity and conservation – GLOBALDIV. *Animal Genetics* **41**, 1-5.
- AL-ATYAT R. M., SALAMEH, N. M. & TABBAA, M. J. (2014). Analysis of genetic diversity and differentiation of sheep populations in Jordan. *Electronic Journal of Biotechnology*.
- ALI B. A. (2003). Genetics similarity among four breeds of sheep in Egypt detected by random amplified polymorphic DNA markers. *African Journal of Biotechnology* **2**, 194-197.
- ALI B. A., HUANG, T.-H., QIN, D.-N. & WANG, X.-M. (2004). A review of random amplified polymorphic DNA (RAPD) markers in fish research. *Reviews in Fish Biology and Fisheries* **14**, 443-453.
- ALLENDORF F. W. (1986). Genetic drift and the loss of alleles versus heterozygosity. *Zoo Biology* **5**, 181-190.
- ALVAREZ I., CAPOTE, J., TRAORÉ, A., FONSECA, N., PÉREZ, K., CUERVO, M., FERNÁNDEZ, I. & GOYACHE, F. (2012). Genetic relationships of the Cuban hair sheep inferred from microsatellite polymorphism. *Small Ruminant Research* **104**, 89-93.
- ALVAREZ I., TRAORÉ, A., TAMBOURA, H. H., KABORE, A., ROYO, L. J., FERNÁNDEZ, I., OUÉDRAOGO-SANOU, G., SAWADOGO, L. & GOYACHE, F. (2009). Microsatellite analysis characterizes Burkina Faso as a

- genetic contact zone between Sahelian and Djallonké sheep. *Animal biotechnology* **20**, 47-57.
- ANDERSON S. (2003). Animal genetic resources and sustainable livelihoods. *Ecological Economics* **45**, 331-339.
- ANDERSON S. & CENTONZE, R. (2007). Property Rights and the Management of Animal Genetic Resources. *World Development* **35**, 1529-1541.
- ANDERSSON L. & GEORGES, M. (2004). Domestic-animal genomics: deciphering the genetics of complex traits. *Nature Reviews Genetics* **5**, 202-212.
- ANDRABI S. M. & MAXWELL, W. M. (2007). A review on reproductive biotechnologies for conservation of endangered mammalian species. *Anim Reprod Sci* **99**, 223-43.
- ASHLEY M. & DOW, B. (1994). The use of microsatellite analysis in population biology: background, methods and potential applications. In *Molecular ecology and evolution: approaches and applications*, pp. 185-201. Springer.
- AVISE J. C., ARNOLD, J., BALL, R. M., BERMINGHAM, E., LAMB, T., NEIGEL, J. E., REEB, C. A. & SAUNDERS, N. C. (1987). Intraspecific Phylogeography: The Mitochondrial DNA Bridge Between Population Genetics and Systematics. *Annual Review of Ecology and Systematics* **18**, 489-522.
- AVISE J. C. & ELLIS, D. (1986). Mitochondrial DNA and the evolutionary genetics of higher animals [and discussion]. *Philosophical Transactions of the Royal Society of London. B, Biological Sciences* **312**, 325-342.
- BANIK S., PANKAJ, P. & NASKAR, S. (2015). Climate Change: Impacts on Livestock Diversity in Tropical Countries. *Livestock Production and Climate Change* **6**, 162.
- BARBATO M., OROZCO-TERWENGEL, P. A., TAPIO, M. & BRUFORD, M. W. (2015). SNeP: a tool to estimate trends in recent effective population size trajectories using genome-wide SNP data. *Frontiers in Genetics* **6**.
- BARENDSE W., VAIMAN, D., KEMP, S., SUGIMOTO, Y., ARMITAGE, S., WILLIAMS, J., SUN, H., EGGEN, A., AGABA, M. & ALEYASIN, S. (1997). A medium-density genetic linkage map of the bovine genome. *Mammalian Genome* **8**, 21-28.
- BARRETT J. C. (1992). *The economic role of cattle in communal farming systems in Zimbabwe*. Overseas Development Institute London.
- BAUMUNG R., CUBRIC-CURIK, V., SCHWEND, K., ACHMANN, R. & SÖLKNER, J. (2006). Genetic characterisation and breed assignment in Austrian sheep breeds using microsatellite marker information. *Journal of Animal Breeding and Genetics* **123**, 265-271.
- BECKMANN J. & SOLLER, M. (1983). Restriction fragment length polymorphisms in genetic improvement: methodologies, mapping and costs. *Theoretical and Applied Genetics* **67**, 35-43.
- BECKMANN J. & SOLLER, M. (1986). Restriction fragment length polymorphisms and genetic improvement of agricultural species. *Euphytica* **35**, 111-124.
- BIRTEEB P. T., PETERS, S. O., YAKUBU, A., ADELEKE, M. A. & OZOJE, M. O. (2012). Multivariate characterisation of the phenotypic traits of Djallonke and Sahel sheep in Northern Ghana. *Tropical animal health and production* **45**, 267-274.
- BISCARINI F., NICOLAZZI, E., STELLA, A., BOETTCHER, P. & GANDINI, G. (2015). Challenges and opportunities in genetic improvement of local livestock breeds. *Frontiers in Genetics* **6**.
- BLEARS M., DE GRANDIS, S., LEE, H. & TREVORS, J. (1998). Amplified fragment length polymorphism (AFLP): a review of the procedure and its applications. *Journal of Industrial Microbiology and Biotechnology* **21**, 99-114.
- BOETTCHER P. J., TIXIER-BOICHARD, M., TORO, M. A., SIMIANER, H., EDING, H., GANDINI, G., JOOST, S., GARCIA, D., COLLI, L., AJMONE-MARSAN, P. & AND THE, G. C. (2010). Objectives, criteria and methods for using molecular genetic data in priority setting for conservation of animal genetic resources. *Animal Genetics* **41**, 64-77.
- BOWLES D., CARSON, A. & ISAAC, P. (2014). Genetic Distinctiveness of the Herdwick Sheep Breed and Two Other Locally Adapted Hill Breeds of the UK: e87823. *PLoS ONE* **9**.

- BRINKS J., CLARK, R., KIEFFER, N. & URICK, J. (1964). Estimates of genetic, environmental and phenotypic parameters in range Hereford females. *Journal of Animal Science* **23**, 711-716.
- BROXHAM E. T. (2015). A case study on strains of Buša cattle structured into a metapopulation to show the potential for use of single-nucleotide polymorphism genotyping in the management of small, cross-border populations of livestock breeds and varieties. *Frontiers in Genetics* **6**.
- BRUFORD M. W., BRADLEY, D. G. & LUIKART, G. (2003). DNA markers reveal the complexity of livestock domestication. *Nature Reviews Genetics* **4**, 900-910.
- BRUFORD M. W. & WAYNE, R. K. (1993). Microsatellites and their application to population genetic studies. *Current Opinion in Genetics & Development* **3**, 939-943.
- BRUNS E. & GLODEK, P. (1999). Animal genetic biodiversity-methodological considerations. In *Deutscher Tropentag*, vol. 14, pp. 1999.
- BUCHANAN F. C., ADAMS, L. J., LITTLEJOHN, R. P., MADDOX, J. F. & CRAWFORD, A. M. (1994). Determination of evolutionary relationships among sheep breeds using microsatellites. *Genomics* **22**, 397-403.
- CASAS E., SHACKELFORD, S., KEELE, J., STONE, R., KAPPES, S. & KOOHARAIE, M. (2000). Quantitative trait loci affecting growth and carcass composition of cattle segregating alternate forms of myostatin. *Journal of Animal Science* **78**, 560-569.
- CHAKRABORTY R. & NEI, M. (1977). Bottleneck Effects on Average Heterozygosity and Genetic Distance with the Stepwise Mutation Model. *Evolution* **31**, 347-356.
- CHEN A., ZHANG, R. & YU, S. (2008). Comparative results of survival of vitrified biopsied goat embryos and mouse morulae. *Turkish Journal of Veterinary and Animal Sciences* **32**, 93-97.
- CHENYAMBUGA S., HANOTTE, O., HIRBO, J., WATTS, P., KEMP, S., KIFARO, G., GWAKISA, P., PETERSEN, P. & REGE, J. (2004). Genetic characterization of indigenous goats of sub-Saharan Africa using microsatellite DNA markers.
- CHESSER R. K., SMITH, M. H. & BRISBIN, I. L. (1980). Management and maintenance of genetic variability in endangered species. *International Zoo Yearbook* **20**, 146-154.
- CHISTIAKOV D. A., HELLEMANS, B. & VOLCKAERT, F. A. M. (2006). Microsatellites and their genomic distribution, evolution, function and applications: A review with special reference to fish genetics. *Aquaculture* **255**, 1-29.
- CLARKE S. M., HENRY, H. M., DODDS, K. G., JOWETT, T. W. D., MANLEY, T. R., ANDERSON, R. M. & MCEWAN, J. C. (2014). A High Throughput Single Nucleotide Polymorphism Multiplex Assay for Parentage Assignment in New Zealand Sheep. *PLoS ONE* **9**, e93392.
- DAVIES G., STEAR, M., BENOITHMAN, M., ABUAGOB, O., KERR, A., MITCHELL, S. & BISHOP, S. (2006). Quantitative trait loci associated with parasitic infection in Scottish blackface sheep. *Heredity* **96**, 252-258.
- DE SOUZA J. M., BATISTA, R. I., MELO, L. M. & FREITAS, V. J. D. F. (2011). Reproductive biotechnologies applied to the conservation of endangered ruminant-past, present and future *Revista Portuguesa de Ciências Veterinárias* **106**, 31-38.
- DIB C., FAURÉ, S., FIZAMES, C., SAMSON, D., DROUOT, N., VIGNAL, A., MILLASSEAU, P., MARC, S., KAZAN, J. & SEBOUN, E. (1996). A comprehensive genetic map of the human genome based on 5,264 microsatellites. *Nature* **380**, 152-154.
- DLUGOSCH K. M. & PARKER, I. M. (2008). Founding events in species invasions: genetic variation, adaptive evolution, and the role of multiple introductions. *Molecular ecology* **17**, 431-449.
- DOBNEY K. & LARSON, G. (2006). Genetics and animal domestication: new windows on an elusive process. *Journal of Zoology* **269**, 261-271.
- DODGSON J. B., CHENG, H. H. & OKIMOTO, R. (1997). DNA marker technology: a revolution in animal genetics. *Poultry science* **76**, 1108-1114.
- DRESZER T. R., KAROLCHIK, D., ZWEIG, A. S., HINRICHS, A. S., RANEY, B. J., KUHN, R. M., MEYER, L. R., WONG, M., SLOAN, C. A. & ROSENBLUM, K. R. (2012). The UCSC Genome Browser database: extensions and updates 2011. *Nucleic acids research* **40**, D918-D923.

- DRUCKER A. G., GOMEZ, V. & ANDERSON, S. (2001). The economic valuation of farm animal genetic resources: a survey of available methods. *Ecological Economics* **36**, 1-18.
- EDING H. & MEUWISSEN, T. (2001). Marker-based estimates of between and within population kinships for the conservation of genetic diversity. *Journal of Animal Breeding and Genetics* **118**, 141-159.
- ENGEL S. R., LINN, R. A., TAYLOR, J. F. & DAVIS, S. K. (1996). Conservation of microsatellite loci across species of artiodactyls: implications for population studies. *Journal of Mammalogy*, 504-518.
- FAN H. & CHU, J.-Y. (2007). A Brief Review of Short Tandem Repeat Mutation. *Genomics, Proteomics & Bioinformatics* **5**, 7-14.
- FAO. (2000). Watch list of Domestic Animal Diversity. FAO and the United Nations' Environmental Programme, Rome, Italy.
- FAO. (2007). *Global Plan of Action for Animal Genetic Resources and the Interlaken Declaration: Adopted by the International Technical Conference on Animal Genetic Resources for Food and Agriculture, Interlaken, Switzerland, 3-7 September 2007*. Commission On Genetic Resources for Food and Agriculture, Food and Agriculture Organization of the United Nations.
- FAO. (2013). Status and Trends of Animal Genetic Resources 2012', pp. 24. Food and Agriculture Organization, Commission on Animal Genetic Resources, Rome, Italy
- FELIUS M., THEUNISSEN, B. & LENSTRA, J. A. (2014). Conservation of cattle genetic resources: the role of breeds. *The Journal of Agricultural Science FirstView*, 1-11.
- FERGUSON A., TAGGART, J., PRODÖHL, P., MCMEEL, O., THOMPSON, C., STONE, C., MCGINNITY, P. & HYNES, R. (1995). The application of molecular markers to the study and conservation of fish populations, with special reference to *Salmo*. *Journal of Fish Biology* **47**, 103-126.
- FRANKHAM R. (1995). Conservation genetics. *Annual review of genetics* **29**, 305-327.
- GARNETT T. (2009). Livestock-related greenhouse gas emissions: impacts and options for policy makers. *Environmental Science & Policy* **12**, 491-503.
- GAUTIER M., FOUCAUD, J., GHARBI, K., CÉZARD, T., GALAN, M., LOISEAU, A., THOMSON, M., PUDLO, P., KERDELHUÉ, C. & ESTOUP, A. (2013). Estimation of population allele frequencies from next-generation sequencing data: pool-versus individual-based genotyping. *Molecular ecology* **22**, 3766-3779.
- GAVRILOVA O., JONAS, M., ERB, K. & HABERL, H. (2010). International trade and Austria's livestock system: Direct and hidden carbon emission flows associated with production and consumption of products. *Ecological Economics* **69**, 920-929.
- GLOBALDIV CONSORTIUM. (2010). Current strategies to Characterize Farm Animal Genetic Resources, pp. 12. EAAP.
- GODDARD M. E. & HAYES, B. J. (2009). Mapping genes for complex traits in domestic animals and their use in breeding programmes. *Nature Reviews Genetics* **10**, 381-391.
- GOE M. & STRANZINGER, G. (2002). Livestock as a component of disaster risk management: quantifying the effect of emergency relief operations on animal genetic resources. *Concept Note prepared for FAO Animal Genetic Resources Group, Rome. Breeding Biology Group, Institute of Animal Sciences, Swiss Federal Institute of Technology, Zurich*.
- GROENEN M. A., MEGENS, H.-J., ZARE, Y., WARREN, W. C., HILLIER, L. W., CROOIJMANS, R. P., VEREIJKEN, A., OKIMOTO, R., MUIR, W. M. & CHENG, H. H. (2011). The development and characterization of a 60K SNP chip for chicken. *BMC Genomics* **12**.
- GROENEVELD L. F., LENSTRA, J. A., EDING, H., TORO, M. A., SCHERF, B., PILLING, D., NEGRINI, R., FINLAY, E. K., JIANLIN, H., GROENEVELD, E., WEIGEND, S. & THE, G. C. (2010). Genetic diversity in farm animals – a review. *Animal Genetics* **41**, 6-31.
- GUO S., SAVOLAINEN, P., SU, J., ZHANG, Q., QI, D., ZHOU, J., ZHONG, Y., ZHAO, X. & LIU, J. (2006). Origin of mitochondrial DNA diversity of domestic yaks. *BMC Evolutionary Biology* **6**, 73.
- GWAKISA P., KEMP, S. & TEALE, A. (1994). Characterization of Zebu cattle breeds in Tanzania using random amplified polymorphic DNA markers. *Animal Genetics* **25**, 89-94.

- HAMBLIN M. T., WARBURTON, M. L. & BUCKLER, E. S. (2007). Empirical comparison of simple sequence repeats and single nucleotide polymorphisms in assessment of maize diversity and relatedness. *PLoS ONE* **2**, e1367.
- HAMMOND K. (1994). Conservation of domestic animal diversity: Global overview. In *Proceedings of the World Congress on Genetics Applied to Livestock Production*, vol. 21, pp. 610.
- HAMMOND K. (1999). Management of farm animal genetic resources. *Economic valuation of animal genetic resources*, 34.
- HANKS J. (2001). Conservation strategies for Africa's large mammals. *Reproduction, fertility and development*.
- HARRISON R. G. (1989). Animal mitochondrial DNA as a genetic marker in population and evolutionary biology. *Trends in Ecology & Evolution* **4**, 6-11.
- HELYAR S. J., HEMMER-HANSEN, J., BEKKEVOLD, D., TAYLOR, M. I., OGDEN, R., LIMBORG, M. T., CARIANI, A., MAES, G. E., DIOPERE, E., CARVALHO, G. R. & NIELSEN, E. E. (2011). Application of SNPs for population genetics of nonmodel organisms: new opportunities and challenges. *Molecular Ecology Resources* **11**, 123-136.
- HENSON E. L. (1992). In situ conservation of livestock and poultry.
- HERRERO-MEDRANO J. M., MEGENS, H.-J., GROENEN, M. A., RAMIS, G., BOSSE, M., PÉREZ-ENCISO, M. & CROOIJMANS, R. P. (2013). Conservation genomic analysis of domestic and wild pig populations from the Iberian Peninsula. *BMC Genetics* **14**, 106.
- HERRERO M., THORNTON, P. K., GERBER, P. & REID, R. S. (2009). Livestock, livelihoods and the environment: understanding the trade-offs. *Current Opinion in Environmental Sustainability* **1**, 111-120.
- HIEMSTRA S. J., DRUCKER, A. G., TVEDT, M. W., LOUWAARS, N., OLDENBROEK, J., AWGICHEW, K., KEBEDE, S. A., BHAT, P. & DA SILVA MARIANTE, A. (2006a). Exchange, use and conservation of animal genetic resources. *Food and Agriculture Organization of the United Nations, Rome*.
- HIEMSTRA S. J., VAN DER LENDE, T. & WOELDERS, H. (2006b). The potential of cryopreservation and reproductive technologies for animal genetic resources conservation strategies. *The role of biotechnology in exploring and protecting agricultural genetic resources. FAO, Rome*, 45-59.
- HOFFMANN I. (2010). Climate change and the characterization, breeding and conservation of animal genetic resources. *Animal Genetics* **41**, 32-46.
- HOFFMANN I. & SCHERF, B. (2005). Animal genetic resources-Time to worry? in Livestock Report 2006 Global perspective, pp. 57-76, Rome Italy.
- HOLT W. V. & PICKARD, A. R. (1999). Role of reproductive technologies and genetic resource banks in animal conservation. *Reviews of reproduction* **4**, 143-150.
- HU Z.-L., PARK, C. A., WU, X.-L. & REECY, J. M. (2013). Animal QTLdb: an improved database tool for livestock animal QTL/association data dissemination in the post-genome era. *Nucleic acids research* **41**, D871-D879.
- HUNDERTMARK K. & VAN DAELE, L. (2010). Founder effect and bottleneck signatures in an introduced, insular population of elk. *Conservation genetics* **11**, 139-147.
- ILRI. (2009). Better Lives Through Livetsock, African Cattle Protected from Killer Disease International Livestock Research Institute.
- JENSEN P. (2006). Domestication—From behaviour to genes and back again. *Applied Animal Behaviour Science* **97**, 3-15.
- KADWELL M., FERNANDEZ, M., STANLEY, H. F., BALDI, R., WHEELER, J. C., ROSADIO, R. & BRUFORD, M. W. (2001). Genetic analysis reveals the wild ancestors of the llama and the alpaca. *Proceedings of the Royal Society of London. Series B: Biological Sciences* **268**, 2575-2584.
- KAHN L. H. (2006). Confronting zoonoses, linking human and veterinary medicine. *Emerging infectious diseases* **12**, 556.
- KANTANEN J., LØVENDAHL, P., STRANDBERG, E., EYTHORSDDOTTIR, E., LI, M.-H., KETTUNEN-PRAEBEL, A., BERG, P. & MEUWISSEN, T. (2015). Utilization of farm animal genetic resources in a changing agro-ecological environment in the Nordic countries. *Frontiers in Genetics* **6**.

- KIJAS J. W., PORTO-NETO, L., DOMINIK, S., REVERTER, A., BUNCH, R., MCCULLOCH, R., HAYES, B. J., BRAUNING, R. & MCEWAN, J. (2014). Linkage disequilibrium over short physical distances measured in sheep using a high-density SNP chip. *Animal Genetics* **45**, 754-757.
- KIJAS J. W., TOWNLEY, D., DALRYMPLE, B. P., HEATON, M. P., MADDOX, J. F., MCGRATH, A., WILSON, P., INGERSOLL, R. G., MCCULLOCH, R. & MCWILLIAM, S. (2009). A genome wide survey of SNP variation reveals the genetic structure of sheep breeds. *PLoS ONE* **4**, e4668.
- KIM S. Y., LI, Y., GUO, Y., LI, R., HOLMKVIST, J., HANSEN, T., PEDERSEN, O., WANG, J. & NIELSEN, R. (2010). Design of association studies with pooled or un-pooled next-generation sequencing data. *Genetic epidemiology* **34**, 479-491.
- KOCHER T. D., THOMAS, W. K., MEYER, A., EDWARDS, S. V., PÄÄBO, S., VILLABLANCA, F. X. & WILSON, A. C. (1989). Dynamics of mitochondrial DNA evolution in animals: amplification and sequencing with conserved primers. *Proceedings of the National Academy of Sciences* **86**, 6196-6200.
- KOEHLER-ROLLEFSON I. & MEYER, H. (2014). Access and Benefit-sharing of Animal Genetic Resources Using the Nagoya Protocol as a Framework for the Conservation and Sustainable Use of Locally.
- KOFER R., NOLTE, V. & SCHLÖTTERER, C. (2015). The impact of library preparation protocols on the consistency of allele frequency estimates in Pool-Seq data. *Molecular Ecology Resources*.
- KOH M. C., LIM, C. H., CHUA, S. B., CHEW, S. T. & PHANG, S. T. W. (1998). Random amplified polymorphic DNA (RAPD) fingerprints for identification of red meat animal species. *Meat Science* **48**, 275-285.
- KÖHLER-ROLLEFSON I. (1997). Indigenous practices of animal genetic resource management and their relevance for the conservation of domestic animal diversity in developing countries. *Journal of Animal Breeding and Genetics* **114**, 231-238.
- KOHN M. H., MURPHY, W. J., OSTRANDER, E. A. & WAYNE, R. K. (2006). Genomics and conservation genetics. *Trends in Ecology & Evolution* **21**, 629-637.
- KOSGEY I. & OKEYO, A. (2007). Genetic improvement of small ruminants in low-input, smallholder production systems: Technical and infrastructural issues. *Small Ruminant Research* **70**, 76-88.
- LACY R. C. (1997). Importance of genetic variation to the viability of mammalian populations. *Journal of Mammalogy*, 320-335.
- LAWRENCE P. R. & PEARSON, R. A. (2002). Use of draught animal power on small mixed farms in Asia. *Agricultural Systems* **71**, 99-110.
- LEVIN I., CRITTENDEN, L. B. & DODGSON, J. B. (1993). Genetic map of the chicken Z chromosome using random amplified polymorphic DNA (RAPD) markers. *Genomics* **16**, 224-230.
- LIU Z. & CORDES, J. (2004). DNA marker technologies and their applications in aquaculture genetics. *Aquaculture* **238**, 1-37.
- LOPES M. S., MENDONÇA, D., ROJER, H., CABRAL, V., BETTENCOURT, S. X. & DA CÂMARA MACHADO, A. (2015). Morphological and genetic characterization of an emerging Azorean horse breed: the Terceira Pony. *Frontiers in Genetics* **6**.
- LUIKART G. & CORNUET, J.-M. (1998). Empirical Evaluation of a Test for Identifying Recently Bottlenecked Populations from Allele Frequency Data. *Conservation Biology* **12**, 228-237.
- LUIKART G., ENGLAND, P. R., TALLMON, D., JORDAN, S. & TABERLET, P. (2003). The Power and Promise of Population Genomics: from Genotyping to Genome Typing. *Nature Reviews: Genetics* **4**, 981-994.
- LUIKART G., FERNÁNDEZ, H., MASHKOUR, M., ENGLAND, P., TABERLET, P., ZEDER, M., EMSHWILLER, E., SMITH, B. & BRADLEY, D. (2006). Origins and diffusion of domestic goats inferred from DNA markers. *Documenting domestication: New genetic and archaeological paradigms*, 294-305.
- MADDOX J. F., DAVIES, K. P., CRAWFORD, A. M., HULME, D. J., VAIMAN, D., CRIBIU, E. P., FREKING, B. A., BEH, K. J., COCKETT, N. E. & KANG, N. (2001). An enhanced linkage map of the sheep genome comprising more than 1000 loci. *Genome research* **11**, 1275-1289.

- MARA L., CASU, S., CARTA, A. & DATTENA, M. (2013). Cryobanking of farm animal gametes and embryos as a means of conserving livestock genetics. *Anim Reprod Sci* **138**, 25-38.
- MARLETTA D., TUPAC-YUPANQUI, I., BORDONARO, S., GARCIA, D., GUASTELLA, A., CRISCIONE, A., CAÑON, J. & DUNNER, S. (2006). Analysis of genetic diversity and the determination of relationships among western Mediterranean horse breeds using microsatellite markers. *Journal of Animal Breeding and Genetics* **123**, 315-325.
- MARTÍN-BURRIEL I., GARCÍA-MURO, E. & ZARAGOZA, P. (1999). Genetic diversity analysis of six Spanish native cattle breeds using microsatellites. *Animal Genetics* **30**, 177-182.
- MATUKUMALLI L. K., LAWLEY, C. T., SCHNABEL, R. D., TAYLOR, J. F., ALLAN, M. F., HEATON, M. P., O'CONNELL, J., MOORE, S. S., SMITH, T. P. & SONSTEGARD, T. S. (2009). Development and characterization of a high density SNP genotyping assay for cattle. *PLoS ONE* **4**, e5350.
- MCCORKLE C. M. & JAMES, V. (1996). The roles of animals in cultural, social, and agro-economic systems. *Sustainable development in Third World countries: applied and theoretical perspectives.*, 25-43.
- McMAHON B. J., TEELING, E. C. & HÖGLUND, J. (2014). How and why should we implement genomics into conservation? *Evolutionary Applications* **7**, 999-1007.
- McMICHAEL A. J., POWLES, J. W., BUTLER, C. D. & UAUY, R. (2007). Food, livestock production, energy, climate change, and health. *The Lancet* **370**, 1253-1263.
- MEDUGORAC I., VEIT-KENSCH, C. E., RAMLJAK, J., BRKA, M., MARKOVIĆ, B., STOJANOVIĆ, S., BYTYQI, H., KOCHOSKI, L., KUME, K. & GRÜNENFELDER, H. P. (2011). Conservation priorities of genetic diversity in domesticated metapopulations: a study in taurine cattle breeds. *Ecology and evolution* **1**, 408-420.
- MENDELSON R. (2003). The challenge of conserving indigenous domesticated animals. *Ecological Economics* **45**, 501-510.
- MIGNON-GRASTEAU S., BOISSY, A., BOUIX, J., FAURE, J.-M., FISHER, A. D., HINCH, G. N., JENSEN, P., LE NEINDRE, P., MORMÈDE, P., PRUNET, P., VANDEPUTTE, M. & BEAUMONT, C. (2005). Genetics of adaptation and domestication in livestock. *Livestock Production Science* **93**, 3-14.
- MORIN P. A., LUIKART, G. & WAYNE, R. K. (2004). SNPs in ecology, evolution and conservation. *Trends in Ecology & Evolution* **19**, 208-216.
- MORRIS J. P. (2006). *The Ethics of Biotechnology*. Chelsea House Publishers, Philadelphia.
- MUNRO R. & ADAMS, D. (1991). Genetic Diversity and Sustainable Agriculture—Implications for Animal Production Systems. *Buffalo and Goats in Asia: Genetic Diversity and its Application, Tulloh, NM (Ed.). Australian Centre for International Agricultural Research (ACIAR), Australia, ISBN-13: 9781863200349*, 123-128.
- NARDONE A., RONCHI, B., LACETERA, N., RANIERI, M. S. & BERNABUCCI, U. (2010). Effects of climate changes on animal production and sustainability of livestock systems. *Livestock Science* **130**, 57-69.
- NASKAR S., GOWANE, G. & CHOPRA, A. (2015). Strategies to Improve Livestock Genetic Resources to Counter Climate Change Impact. In *Climate Change Impact on Livestock: Adaptation and Mitigation* (ed. V. Sejian, J. Gaughan, L. Baumgard and C. Prasad), pp. 441-475. Springer India.
- NEI M., MARUYAMA, T. & CHAKRABORTY, R. (1975). The bottleneck effect and genetic variability in populations. *Evolution*, 1-10.
- NEWMAN D. & PILSON, D. (1997). Increased Probability of Extinction Due to Decreased Genetic Effective Population Size: Experimental Populations of *Clarkia pulchella*. *Evolution* **51**, 354-362.
- NOTTER D., MARIANTE, A. D. S. & SHENG, Z. (1994). Modern approaches to active conservation of domestic animal diversity. In *Proc. 5th World Congress on Genetic Applied to Livestock Production. Dept. Animal and Poultry Science, University of Guelph. Ontario, Canada*, vol. 21, pp. 509-516.

- O'BRIEN J. K., STEINMAN, K. J. & ROBECK, T. R. (2009). Application of sperm sorting and associated reproductive technology for wildlife management and conservation. *Theriogenology* **71**, 98-107.
- OLSON H., BETTON, G., ROBINSON, D., THOMAS, K., MONRO, A., KOLAJA, G., LILLY, P., SANDERS, J., SIPES, G. & BRACKEN, W. (2000). Concordance of the toxicity of pharmaceuticals in humans and in animals. *Regulatory Toxicology and Pharmacology* **32**, 56-67.
- PAIVA S., FARIA, D., SILVÉRIO, V., MCMANUS, C., EGITO, A., DERGAM, J., GUIMARÃES, S., CASTRO, S., ALBUQUERQUE, M. & MARIANTE, A. (2005). Genetic variability among Brazilian sheep using microsatellites. *Proc. The Role of biotechnology, Turin, March*, 5-7.
- PARSONS Y. M. & SHAW, K. (2001). Species boundaries and genetic diversity among Hawaiian crickets of the genus *Laupala* identified using amplified fragment length polymorphism. *Molecular ecology* **10**, 1765-1772.
- PEPIN L. & NGUYEN, T. (1994). Blood groups and protein polymorphisms in five goat breeds (*Capra hircus*). *Animal Genetics* **25**, 333-336.
- PHILIPSSON J., REGE, J. & OKEYO, A. (2006). Sustainable breeding programmes for tropical farming systems. *Animal genetics training resource, Version 2*, 2006.
- PHILIPSSON J., ZONABEND, E., BETT, R. & OKEYO, A. (2011). Swedish University of Agricultural Sciences (SLU), Department of Animal Breeding and Genetics, PO Box 7023, S-750 07 Uppsala, Sweden. 2 International Livestock Research Institute (ILRI), PO Box 30709, Nairobi 00100, Kenya.
- PINTADO B. & HOURCADE, J. (2011). Cryopreservation. In *Advanced Protocols for Animal Transgenesis*, pp. 577-599. Springer.
- PRENTICE J. R. & ANZAR, M. (2011). Cryopreservation of Mammalian Oocyte for Conservation of Animal Genetics. *Veterinary Medicine International* **2011**, 11.
- PRICE E. O. (1999). Behavioral development in animals undergoing domestication. *Applied Animal Behaviour Science* **65**, 245-271.
- QANBARI S., PAUSCH, H., JANSEN, S., SOMEL, M., STROM, T. M., FRIES, R., NIELSEN, R. & SIMIANER, H. (2014). Classic selective sweeps revealed by massive sequencing in cattle. *PLoS genetics* **10**, e1004148.
- RAMSTAD K. M., WOODY, C. A., SAGE, G. K. & ALLENDORF, F. W. (2004). Founding events influence genetic population structure of sockeye salmon (*Oncorhynchus nerka*) in Lake Clark, Alaska. *Molecular ecology* **13**, 277-290.
- REGE J. E. O. (2003). Defining livestock breeds in the context of community-based management of farm animal genetic resources. *Community-based management of animal genetic resources*, 27-36.
- REGE J. E. O. & GIBSON, J. P. (2003). Animal genetic resources and economic development: issues in relation to economic valuation. *Ecological Economics* **45**, 319-330.
- REGE J. E. O., MARSHALL, K., NOTENBAERT, A., OJANGO, J. M. K. & OKEYO, A. M. (2011). Pro-poor animal improvement and breeding — What can science do? *Livestock Science* **136**, 15-28.
- REVERTER A., JOHNSTON, D., FERGUSON, D., PERRY, D., GODDARD, M., BURROW, H., ODDY, V., THOMPSON, J. & BINDON, B. (2003). Genetic and phenotypic characterisation of animal, carcass, and meat quality traits from temperate and tropically adapted beef breeds. 4. Correlations among animal, carcass, and meat quality traits. *Crop and Pasture Science* **54**, 149-158.
- RISCHKOWSKY B., PILLING, D., FOOD, C. O. G. R. F. & AGRICULTURE. (2007). *The State of the World's Animal Genetic Resources for Food and Agriculture*. Commission on Genetic Resources for Food and Agriculture, Food and Agriculture Organization of the United Nations.
- ROHRER G. A., ALEXANDER, L. J., KEELE, J. W., SMITH, T. P. & BEATTIE, C. W. (1994). A microsatellite linkage map of the porcine genome. *Genetics* **136**, 231-245.
- ROHRER G. A. & KEELE, J. (1998). Identification of quantitative trait loci affecting carcass composition in swine: I. Fat deposition traits. *Journal of Animal Science* **76**, 2247-2254.

- RUANE J. & SONNINO, A. (2011). Agricultural biotechnologies in developing countries and their possible contribution to food security. *Journal of biotechnology* **156**, 356-363.
- RUSO V., MARTELLI, M., MAURO, A., DI GIACINTO, O., NARDINOCCHI, D. & BERARDINELLI, P. (2007). Nuclear remodelling in growing oocytes of sheep. *Veterinary research communications* **31**, 201-204.
- SANTOS R., AFFONSO, P., DINIZ, D., MEDRADO, A., SILVA, K. & CARNEIRO, P. (2013). Identification of naturalized goat breeds under conservation from northeastern Brazil using chromosomal markers. *Genetics and Molecular Research* **12**, 3193-3200.
- SAVELKOUL P., AARTS, H., DE HAAS, J., DIJKSHOORN, L., DUIM, B., OTSEN, M., RADEMAKER, J., SCHOULS, L. & LENSTRA, J. (1999). Amplified-fragment length polymorphism analysis: the state of an art. *Journal of clinical microbiology* **37**, 3083-3091.
- SAYERS E. W., BARRETT, T., BENSON, D. A., BOLTON, E., BRYANT, S. H., CANESE, K., CHETVERNIN, V., CHURCH, D. M., DICUCCIO, M. & FEDERHEN, S. (2011). Database resources of the national center for biotechnology information. *Nucleic acids research* **39**, D38-D51.
- SCHLÖTTERER C., KOFLER, R., VERSACE, E., TOBLER, R. & FRANSSSEN, S. (2014). Combining experimental evolution with next-generation sequencing: a powerful tool to study adaptation from standing genetic variation. *Heredity*.
- SHAND H. (1997). *Human Nature : Agricultural Biodiversity and Farm Based Food Food security*, Ottawa : Rural Advancement Foundation International.
- SHIELDS S. & ORME-EVANS, G. (2015). The Impacts of Climate Change Mitigation Strategies on Animal Welfare. *Animals* **5**, 361-394.
- SHIOMI M., ITO, T., YAMADA, S., KAWASHIMA, S. & FAN, J. (2003). Development of an animal model for spontaneous myocardial infarction (WHHLMI rabbit). *Arteriosclerosis, thrombosis, and vascular biology* **23**, 1239-1244.
- SIGNORELLO G. & PAPPALARDO, G. (2003). Domestic animal biodiversity conservation: a case study of rural development plans in the European Union. *Ecological Economics* **45**, 487-499.
- SIMON D. (1999). European approaches to conservation of farm animal genetic resources. *Animal Genetic Resources Information* **25**, 77-97.
- SUNNUCKS P. (2000). Efficient genetic markers for population biology. *Trends in Ecology & Evolution* **15**, 199-203.
- TAKEZAKI N. & NEI, M. (1996). Genetic distances and reconstruction of phylogenetic trees from microsatellite DNA. *Genetics* **144**, 389-399.
- TAKEZAKI N. & NEI, M. (2008). Empirical tests of the reliability of phylogenetic trees constructed with microsatellite DNA. *Genetics* **178**, 385-392.
- TEMPLETON A. R. (1980). The theory of speciation via the founder principle. *Genetics* **94**, 1011-1038.
- TEWELDMEHIDIN M. Y. & CONROY, A. (2010). The economic importance of draught oxen on small farms in Namibia's Eastern Caprivi Region. *African Journal of Agricultural Research* **5**, 928-934.
- THORNTON P. K., BOONE, R. B., GALVIN, K. A., BURNSILVER, S. B., WAITHAKA, M. M., KUYIAH, J., KARANJA, S., GONZÁLEZ-ESTRADA, E. & HERRERO, M. (2007). Coping strategies in livestock-dependent households in east and southern Africa: a synthesis of four case studies. *Human Ecology* **35**, 461-476.
- THORNTON P. K., VAN DE STEEG, J., NOTENBAERT, A. & HERRERO, M. (2009). The impacts of climate change on livestock and livestock systems in developing countries: A review of what we know and what we need to know. *Agricultural Systems* **101**, 113-127.
- THURSTON L. M., SIGGINS, K., MILEHAM, A. J., WATSON, P. F. & HOLT, W. V. (2002). Identification of amplified restriction fragment length polymorphism markers linked to genes controlling boar sperm viability following cryopreservation. *Biology of Reproduction* **66**, 545-554.
- TISDELL C. (2003). Socioeconomic causes of loss of animal genetic diversity: analysis and assessment. *Ecological Economics* **45**, 365-376.
- TIXIER-BOICHARD M. (2014). Status and gaps in characterization of animal genetic resources. In *10th World Congress on Genetics Applied to Livestock Production*. Asas.

- TORO M. A., FERNÁNDEZ, J. & CABALLERO, A. (2009). Molecular characterization of breeds and its use in conservation. *Livestock Science* **120**, 174-195.
- UNITED NATIONS. (2000). Millennium Development Goals and Beyond 2015.
- VANHALA T., TUISKULA-HAAVISTO, M., ELO, K., VILKKI, J. & MAKI-TANILA, A. (1998). Evaluation of genetic variability and genetic distances between eight chicken lines using microsatellite markers. *Poultry science* **77**, 783-790.
- VASCONCELLOS L. P. D. M. K., TAMBASCO-TALHARI, D., PEREIRA, A. P., COUTINHO, L. L. & REGITANO, L. C. D. A. (2003). Genetic characterization of Aberdeen Angus cattle using molecular markers. *Genetics and Molecular Biology* **26**, 133-137.
- VIGNAL A., MILAN, D., SANCRISTOBAL, M. & EGGEN, A. (2002). A review on SNP and other types of molecular markers and their use in animal genetics. *Genetics Selection Evolution* **34**, 275-306.
- VIGNE J.-D. (2011). The origins of animal domestication and husbandry: A major change in the history of humanity and the biosphere. *Comptes Rendus Biologies* **334**, 171-181.
- VISSCHER P., WOOLLIAMS, J., SMITH, D. & WILLIAMS, J. (2002). Estimation of pedigree errors in the UK dairy population using microsatellite markers and the impact on selection. *Journal of Dairy Science* **85**, 2368-2375.
- VODIČKA P., SMETANA, K., DVOŘÁNKOVÁ, B., EMERICK, T., XU, Y. Z., OUREDNIK, J., OUREDNIK, V. & MOTLÍK, J. A. N. (2005). The Miniature Pig as an Animal Model in Biomedical Research. *Annals of the New York Academy of Sciences* **1049**, 161-171.
- WAFULA P., JIANLIN, H., SANGARE, N., SOWE, J., COLY, R., DIALLO, B. & HANOTTE, O. (2005). Genetic characterization of West African Djallonke sheep using microsatellite markers. In *International workshop "The role of biotechnology for the characterization and conservation of crop, forestry, animal and fishery genetic resources"*. Villa Gualino, Turin, pp. 177-178.
- WASKO A. P. & GALETTI, P. M. (2003). PCR primed with minisatellite core sequences yields species-specific patterns and assessment of population variability in fishes of the genus Brycon. *Journal of Applied Ichthyology* **19**, 109-113.
- WITKO-SARSAT V., FRIEDLANDER, M., CAPELLERE-BLANDIN, C., NGUYEN-KHOA, T., NGUYEN, A. T., ZINGRAFF, J., JUNGERS, P. & DESCAMPS-LATSCHA, B. (1996). Advanced oxidation protein products as a novel marker of oxidative stress in uremia. *Kidney international* **49**, 1304-1313.
- WOLLNY C. B. A. (2003). The need to conserve farm animal genetic resources in Africa: should policy makers be concerned? *Ecological Economics* **45**, 341-351.
- WORLEY K., COLLET, J., SPURGIN, L. G., CORNWALLIS, C., PIZZARI, T. & RICHARDSON, D. S. (2010). MHC heterozygosity and survival in red junglefowl. *Molecular ecology* **19**, 3064-3075.
- XIAO-YONG C., SHU-JUN, T., RUN-ZI, S., SHU-CHUN, S., ZHU-JUN, Z., JUN-JIE, L. & WEI-TAO, D. (2008). Induction of lamb follicular development and embryo production in vitro. *Chinese Journal of Agricultural Biotechnology* **5**, 257-262.
- YAPI-GNAORÉ C. V., DAGNOGO, B. & OYA, B. (2003). Community-based livestock improvement and conservation: experiences from open-nucleus breeding programmes in West Africa. *Community-based management of animal genetic resources*, 89.
- YORK R. & MANCUS, P. (2013). The Invisible Animal Anthrozoology and Macrosociology. *Sociological Theory* **31**, 75-91.
- ZANE L., BARGELLONI, L. & PATARNELLO, T. (2002). Strategies for microsatellite isolation: a review. *Molecular ecology* **11**, 1-16.
- ZEDER M. A., EMSHWILLER, E., SMITH, B. D. & BRADLEY, D. G. (2006). Documenting domestication: the intersection of genetics and archaeology. *Trends in Genetics* **22**, 139-155.
- ZHANG D.-X. & HEWITT, G. M. (1996). Nuclear integrations: challenges for mitochondrial DNA markers. *Trends in Ecology & Evolution* **11**, 247-251.
- ZHAO S., ZHENG, P., DONG, S., ZHAN, X., WU, Q., GUO, X., HU, Y., HE, W., ZHANG, S. & FAN, W. (2013). Whole-genome sequencing of giant pandas provides insights into demographic history and local adaptation. *Nature genetics* **45**, 67-71.

